

Gene Section

Mini Review

PLAGL1 (pleomorphic adenoma gene-like 1)

Abbas Abdollahi

Sbarro Institute for Cancer Research and Molecular Medicine, College of Science and Technology, BioLife Sciences Building, Suite 446, Temple University, 1900 North 12th Street, Philadelphia, PA 19122, USA

Published in Atlas Database: August 2007

Online updated version: <http://AtlasGeneticsOncology.org/Genes/PLAGL1ID41737ch6q24.html>

DOI: 10.4267/2042/38498

This work is licensed under a Creative Commons Attribution-Non-commercial-No Derivative Works 2.0 France Licence.

© 2008 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Identity

Hugo: PLAGL1

Other names: DKFZp781P1017; LOT1; MGC126275; MGC126276; ZAC; ZAC1

Location: 6q24.2

DNA/RNA

Description

The genome is about 64 kbp with six exons and five introns. The major mRNA transcript is about 4.7 kb in size.

Transcription

At least three splicing variants (see figure).

Protein

Description

The PLAGL1 protein is a seven C2H2-type zinc finger protein. The seven zinc finger domain is located at the amino terminal region, from the amino acid residue 1 to 210. Additional features of note are proline and glutamine rich regions at the carboxyl terminal portion (residues 220 to 444). There are two nuclear localization signals at the amino terminal region.

Expression


Ovary, breast, brain, liver, spleen, thymus, prostate, uterus, testis, intestine, colon, leukocytes.

Localisation

Nuclear.

Function

The PLAGL1 protein is a candidate tumor suppressor gene and has been shown to have transactivation and



Schematic view of the PLAGL1 gene. A. The six exons, shown in red, are 326; 72; 1443; 75; 475; 2380 bp, respectively. The introns are approximately 39; 2.6; 4.2; 12.5; 5.5 kbp in size, respectively. B. The figure shows three splice variants of PLAGL1; the exons are shown as a box.

Homology

Homologous to the mouse and Rat *plagl1* and to the human PLAG1 and PLAGL2 proteins.

Mutations

Note: Mutation has not been found in the PLAGL1 coding region.

Implicated in

Carcinogenicity

Note: Alteration of the gene expression was found to be a potential genetic event silencing the PLAGL1 gene in cancers such as breast primary tumors, ovarian primary tumors and tumor-derived cell lines, basal cell carcinoma, head and neck squamous cell carcinoma (HNSCC), and extraskeletal myxoid chondrosarcoma (EMC). Several mechanisms have been shown to regulate the PLAGL1 gene expression, including growth factor receptor activation, epigenetic factors, maternal imprinting, and loss of heterozygosity (LOH). Allelic deletion of 6q24-q25, the PLAGL1 locus, in the tumor tissues has been shown in the cancers of ovarian, breast, HNSCC, and pheochromocytomas (PCCs).

Disease

Cancer

Transient neonatal diabetes (TNDM)

Note: Initial reports suggested that transient neonatal diabetes mellitus (TNDM), a rare condition characterized in the patients by intrauterine growth retardation, dehydration, failure to thrive, and hyperglycemia due to a lack of normal insulin secretion, is associated with paternal uniparental disomy (UPD) of chromosome 6 (UPD 6). Later studies showed the involvement of an imprinted gene in this disease within the chromosomal region 6q24.1-q24.3 between markers D6S1699 and D6S1010. Analysis of the CpG islands in the TNDM critical region, using DNA from TNDM patients with paternal UPD 6 and normal controls, suggested PLAGL1/LOT1/ZAC1 as a candidate imprinted gene for this disease.

Disease

Diabetes mellitus

To be noted

Note: PLAGL1 has been implicated in embryonic development. The gene is variably expressed in different tissues and stages during development. Inactivation of this maternally repressed gene resulted in intrauterine growth restriction, altered bone formation, and neonatal lethality.

References

- Abdollahi A, Roberts D, Godwin AK, Schultz DC, Sonoda G, Testa JR, Hamilton TC. Identification of a zinc-finger gene at 6q25: a chromosomal region implicated in development of many solid tumors. *Oncogene* 1997;14(16):1973-1979.
- Abdollahi A, Godwin AK, Miller PD, Getts LA, Schultz DC, Taguchi T, Testa JR, Hamilton TC. Identification of a gene containing zinc-finger motifs based on lost expression in malignant transformed rat ovarian surface epithelial cells. *Cancer Res* 1997;57(10):2029-2034.
- Spengler D, Villalba M, Hoffmann A, Pantaloni C, Houssami S, Bockaert J, Journot L. Regulation of apoptosis and cell cycle arrest by Zac1, a novel zinc finger protein expressed in the pituitary gland and the brain. *EMBO J* 1997;16(10):2814-2825.
- Gillardone F, Hata R, Hossmann KA. Delayed up-regulation of Zac1 and PACAP type I receptor after transient focal cerebral ischemia in mice. *Brain Res Mol Brain Res* 1998;61(1-2):207-210.
- Kas K, Voz ML, Hensen K, Meyen E, Van de Ven WJ. Transcriptional activation capacity of the novel PLAG family of zinc finger proteins. *J Biol Chem* 1998;273(36):23026-23032.
- Varrault A, Ciani E, Apiou F, Bilanges B, Hoffmann A, Pantaloni C, Bockaert J, Spengler D, Journot L. hZAC encodes a zinc finger protein with antiproliferative properties and maps to a chromosomal region frequently lost in cancer. *Proc Natl Acad Sci USA* 1998;95(15):8835-8840.
- Abdollahi A, Bao R, Hamilton TC. LOT1 is a growth suppressor gene down-regulated by the epidermal growth factor receptor ligands and encodes a nuclear zinc-finger protein. *Oncogene* 1999;18(47):6477-6487.
- Bilanges B, Varrault A, Basyuk E, Rodriguez C, Mazumdar A, Pantaloni C, Bockaert J, Theillet C, Spengler D, Journot L. Loss of expression of the candidate tumor suppressor gene ZAC in breast cancer cell lines and primary tumors. *Oncogene* 1999;18(27):3979-3988.
- Pagotto U, Arzberger T, Ciani E, Lezoualc'h F, Pilon C, Journot L, Spengler D, Stalla GK. Inhibition of Zac1, a new gene differentially expressed in the anterior pituitary, increases cell proliferation. *Endocrinology* 1999;140(2):987-996.
- Arima T, Drewell RA, Oshimura M, Wake N, Surani MA. A novel imprinted gene, HYMAI, is located within an imprinted domain on human chromosome 6 containing ZAC. *Genomics* 2000;67(3):248-255.
- Huang SM, Stallcup MR. Mouse Zac1, a transcriptional coactivator and repressor for nuclear receptors. *Mol Cell Biol* 2000;20(5):1855-1867.
- Kamiya M, Judson H, Okazaki Y, Kusakabe M, Muramatsu M, Takada S, Takagi N, Arima T, Wake N, Kamimura K, Satomura K, Hermann R, Bonthron DT, Hayashizaki Y. The cell cycle control gene ZAC/PLAGL1 is imprinted--a strong candidate gene for transient neonatal diabetes. *Hum Mol Genet* 2000;9(3):453-460.
- Pagotto U, Arzberger T, Theodoropoulou M, Grubler Y, Pantaloni C, Saeger W, Losa M, Journot L, Stalla GK, Spengler D. The expression of the antiproliferative gene ZAC is lost or highly reduced in nonfunctioning pituitary adenomas. *Cancer Res* 2000;60(24):6794-6799.
- Piras G, El Kharroubi A, Kozlov S, Escalante-Alcalde D, Hernandez L, Copeland NG, Gilbert DJ, Jenkins NA, Stewart CL. Zac1 (Lot1), a potential tumor suppressor gene, and the gene for epsilon-sarcoglycan are maternally imprinted genes:

identification by a subtractive screen of novel uniparental fibroblast lines. *Mol Cell Biol* 2000;20(9):3308-3315.

Bilanges B, Varrault A, Mazumdar A, Pantaloni C, Hoffmann A, Bockaert J, Spengler D, Journot L. Alternative splicing of the imprinted candidate tumor suppressor gene ZAC regulates its antiproliferative and DNA binding activities. *Oncogene* 2001;20(10):1246-1253.

Huang SM, Schönthal AH, Stallcup MR. Enhancement of p53-dependent gene activation by the transcriptional coactivator Zac1. *Oncogene* 2001;20(17):2134-2143.

Valente T, Auladell C. Expression pattern of Zac1 mouse gene, a new zinc-finger protein that regulates apoptosis and cellular cycle arrest, in both adult brain and along development. *Mech Dev* 2001;108(1-2):207-211.

Varrault A, Bilanges B, Mackay DJ, Basyuk E, Ahr B, Fernandez C, Robinson DO, Bockaert J, Journot L. Characterization of the methylation-sensitive promoter of the imprinted ZAC gene supports its role in transient neonatal diabetes mellitus. *J Biol Chem* 2001;276(22):18653-18656.

Hensen K, Van Valckenborgh IC, Kas K, Van de Ven WJ, Voz ML. The tumorigenic diversity of the three PLAG family members is associated with different DNA binding capacities. *Cancer Res* 2002;62(5):1510-1517.

Mackay DJ, Coupe AM, Shield JP, Storr JN, Temple IK, Robinson DO. Relaxation of imprinted expression of ZAC and HYMAI in a patient with transient neonatal diabetes mellitus. *Hum Genet* 2002;110(2):139-144.

Smith RJ, Arnaud P, Konfortova G, Dean WL, Beechey CV, Kelsey G. The mouse Zac1 locus: basis for imprinting and comparison with human ZAC. *Gene* 2002;292(1-2):101-112.

Abdollahi A, Pisarcik D, Roberts D, Weinstein J, Cairns P, Hamilton TC. LOT1 (PLAGL1/ZAC1), the candidate tumor suppressor gene at chromosome 6q24-25, is epigenetically regulated in cancer. *J Biol Chem* 2003;278(8):6041-6049.

Ciani E, Frenquelli M, Contestabile A. Developmental expression of the cell cycle and apoptosis controlling gene, Lot1, in the rat cerebellum and in cultures of cerebellar granule cells. *Brain Res Dev Brain Res* 2003;142(2):193-202.

Hoffmann A, Ciani E, Boeckardt J, Holsboer F, Journot L, Spengler D. Transcriptional activities of the zinc finger protein Zac are differentially controlled by DNA binding. *Mol Cell Biol* 2003;23(3):988-1003.

Cvetkovic D, Pisarcik D, Lee C, Hamilton TC, Abdollahi A. Altered expression and loss of heterozygosity of the LOT1 gene in ovarian cancer. *Gynecol Oncol* 2004;95(3):449-455.

Koy S, Hauses M, Appelt H, Friedrich K, Schackert HK, Eckelt U. Loss of expression of ZAC/LOT1 in squamous cell carcinomas of head and neck. *Head Neck* 2004;26(4):338-344.

Valente T, Domínguez MI, Bellmann A, Journot L, Ferrer I, Auladell C. Zac1 is up-regulated in neural cells of the limbic system of mouse brain following seizures that provoke strong cell activation. *Neuroscience* 2004;128(2):323-336.

Arima T, Kamikihara T, Hayashida T, Kato K, Inoue T, Shirayoshi Y, Oshimura M, Soejima H, Mukai T, Wake N. ZAC, LIT1 (KCNQ1OT1) and p57KIP2 (CDKN1C) are in an imprinted gene network that may play a role in Beckwith-Wiedemann syndrome. *Nucleic Acids Res* 2005;33(8):2650-2660.

Basyuk E, Coulon V, Le Digarcher A, Coisy-Quivy M, Moles JP, Gandarillas A, Journot L. The candidate tumor suppressor gene ZAC is involved in keratinocyte differentiation and its

expression is lost in basal cell carcinomas. *Mol Cancer Res* 2005;3(9):483-492.

Contestabile A, Fila T, Bartesaghi R, Ciani E. Cyclic AMP-mediated regulation of transcription factor Lot1 expression in cerebellar granule cells. *J Biol Chem* 2005;280(39):33541-33551.

Kamikihara T, Arima T, Kato K, Matsuda T, Kato H, Douchi T, Nagata Y, Nakao M, Wake N. Epigenetic silencing of the imprinted gene ZAC by DNA methylation is an early event in the progression of human ovarian cancer. *Int J Cancer* 2005;115(5):690-700.

Poulin H, Labelle Y. The PLAGL1 gene is down-regulated in human extraskeletal myxoid chondrosarcoma tumors. *Cancer Lett* 2005;227(2):185-191.

Valente T, Junyent F, Auladell C. Zac1 is expressed in progenitor/stem cells of the neuroectoderm and mesoderm during embryogenesis: differential phenotype of the Zac1-expressing cells during development. *Dev Dyn* 2005;233(2):667-679.

Arima T, Yamasaki K, John RM, Kato K, Sakumi K, Nakabeppu Y, Wake N, Kono T. The human HYMAI/PLAGL1 differentially methylated region acts as an imprint control region in mice. *Genomics* 2006;88(5):650-658.

Arima T, Wake N. Establishment of the primary imprint of the HYMAI/PLAGL1 imprint control region during oogenesis. *Cytogenet Genome Res* 2006;113(1-4):247-252.

Barz T, Hoffmann A, Panhuysen M, Spengler D. Peroxisome proliferator-activated receptor gamma is a Zac target gene mediating Zac antiproliferation. *Cancer Res* 2006;66(24):11975-11982.

Hoffmann A, Barz T, Spengler D. Multitasking C2H2 zinc fingers link Zac DNA binding to coordinated regulation of p300-histone acetyltransferase activity. *Mol Cell Biol* 2006;26(14):5544-5557.

Varrault A, Gueydan C, Delalbre A, Bellmann A, Houssami S, Aknin C, Severac D, Chotard L, Kahli M, Le Digarcher A, Pavlidis P, Journot L. Zac1 regulates an imprinted gene network critically involved in the control of embryonic growth. *Dev Cell* 2006;11(5):711-722.

Abdollahi A. LOT1 (ZAC1/PLAGL1) and its family members: mechanisms and functions. *J Cell Physiol* 2007;210(1):16-25. (Review).

Ma L, Cantrup R, Varrault A, Colak D, Klenin N, Gotz M, McFarlane S, Journot L, Schuurmans C. Zac1 functions through TGFbeta11 to negatively regulate cell number in the developing retina. *Neural Develop* 2007;2:11.

Valleley EM, Cordery SF, Bonthron DT. Tissue-specific imprinting of the ZAC/PLAGL1 tumour suppressor gene results from variable utilization of monoallelic and biallelic promoters. *Hum Mol Genet* 2007;16(8):972-981.

Wu MH, Huang CJ, Liu ST, Liu PY, Ho CL, Huang SM. Physical and functional interactions of human papillomavirus E2 protein with nuclear receptor coactivators. *Biochem Biophys Res Commun* 2007;356(3):523-528.

This article should be referenced as such:

Abdollahi A. PLAGL1 (pleomorphic adenoma gene-like 1). *Atlas Genet Cytogenet Oncol Haematol*.2008;12(2):127-129.
